

GENITAL INFECTIONS

Allegato

All-SDF.LAB0	3.	55
--------------	----	----

Pagina 1 di 2

Aggiornato il: 05/07/2024

Fondazione TERESA CAMPLANI BORATORIO ANALISI

ISTRUCTIONS FOR THE PATIENTS

VAGINAL SWAB

Ш	Gential secretions should not be confected during the menstrual period and at least three
	days afterthe end of blood loss.
	Vaginal douches should not be performed and antibiotics should not be administered in
	the 5 daysprior to the sampling.
	Refrain from sexual intercourses within 48 hours prior to the sampling.

STREPTOCOCCUS AGALACTIAE RESEARCH IN VAGINE-RECTAL BUFFER OF **PREGNANTWOMEN**

	Vaginal douches should not be performed and antibiotics should not be administered in
	the 5 daysprior to the sampling.
_	

☐ Refrain from sexual intercourses within 48 hours prior to the sampling.

VULVAR SWAB

- Genital secretions should not be collected during the menstrual period and at least three days afterthe end of blood loss.
- Vaginal douches should not be performed and antibiotics should not be administered in the 5 daysprior to the sampling.
- Refrain from sexual intercourses within 48 hours prior to the sampling.
- Do not wash the affected area within 12 hours prior to the sampling.

ENDOCERVICAL SWAB

- Genital secretions should not be collected during the menstrual period and at least three three daysafter the end of blood loss.
- Antibiotics should not be administered in the 30 days (PCR/NAAT) or 5 days (Neisseria gonorrhoeae culture) prior to the sampling.
- Refrain from sexual intercourses within 48 hours prior to the sampling.

URETHRAL SAMPLING

- Antibiotics should not be administered within 30 days (PCR/NAAT) or 5 days (culture) prior to the sampling.
- Refrain from sexual intercourses within 48 hours prior to the sampling.
- Sampling should be carried out in the morning before urination or, alternatively, two to three hoursafter the last urination.
- Concerning PCR/NAAT test
 - Get a sterile wide mouth container equipped with screw cap suitable for "urinocoltura"



Fondazione
TERESA CAMPLANI
LABORATORIO ANALISI



Allegato

GENITAL INFECTIONS

All-SDF.LAB03.55

Pagina 2 di 2

Aggiornato il: 05/07/2024

(urine culture).

- Write the patient's surname, first name, date of birth or paste the barcode label.
- Collect the first stream of urine in the morning or, alternatively, the second one (at least twoto three hours after the last urination)
- Close carefully the container.